


Commonwealth of Virginia
STATE CORPORATION COMMISSION

DIVISION OF ENERGY REGULATION

MEMORANDUM

May 23, 2016

TO: Document Control Center

FROM: Daisy Maldonado, Energy Regulation 

RE: PUE-2015-00107 –Virginia Electric and Power Company - for Approval and Certification of Electric Facilities Haymarket 230 kV Double Circuit Transmission Line and 230-34.5 kV Haymarket Substation.

SCC-CLERK'S OFFICE
DOCUMENT CONTROL CENTER
2016 MAY 23 P 3:45

160550042

Attached are two (2) case comments submitted to the Commission for consideration. Please enter in the case jacket for Case No. PUE-2015-00107. Thank you.

Mr. Andrew Shropshire, et al.

NOT FOR DISTRIBUTION

169550042

Daisy Maldonado

From: ashropshire77@gmail.com
Sent: Sunday, May 22, 2016 10:13 AM
To: PUE_Comments; Charlotte.p.mcafee@dom.com; vlink@mcguirewoods.com; jvalaika@mcguirewoods.com
Subject: Case Comments Submission for Case # PUE-2015-00107
Importance: High


The following case comments were submitted online Sunday, May 22, 2016 at 10:12:41 AM

Full Name: Mr. Andrew Shropshire
Group or Organization:
Address Line One: 16036 Sheringham Way
Address Line Two:
City, State, Zip: Gainesville, VA 20155
Email: ashropshire77@gmail.com
Case Number: PUE-2015-00107

Comments: SCC and Staff, I would like to bring to your attention that a 63MW fuel cell park was recently approved by Connecticut. For testimony and other document's please see that state's web site at (<http://www.ct.gov/csc/cwp/view.asp?a=2397&q=570126>). This installation is of comparable size to one that could be used to satisfy the power needs in an alternative way to a proposed electric transmission line (by extending the gas transmission line owned by Columbia Gas at the intersection of route 29 and 66). As stated elsewhere, drawing the energy from transmitted gas rather than transmitted electricity is significantly cheaper in terms of fuel costs (electricity costs are more than double natural gas at 47% conversion rates for typical fuel cells). A gas transmission line would also have less visual/negative economic impact on the community than any of the proposed transmission line routes. Fuel cells such as the DFC3000 by Fuel Cell Inc (which are used in the CT application above) or the PureCell 400 by Doosan Fuel Cell America Inc can be aggregated to produce the 100MW or other energy needs, utilizing a small amount of land (as proven by their previous installations of comparable power systems). Dominion Power owns a 15MW plant of 5 DFC3000 fuel cells in Connecticut which it recently built. Dominion executives can be seen here promoting and endorsing the DFC3000 (see <http://www.fuelcellenergy.com/news-resources/videos/> (see upper right video)). Regarding costs: Scott Samuelson, director of the National Fuel Cell Research Center at the University of California, Irvine, quoted in a CNBC report, claims, "The fuel-cell companies that are public are beginning to demonstrate costs that are beginning to approach \$3,000 per kilowatt." (from <http://www.greentechmedia.com/articles/read/Fuel-Cells-2016-Within-Striking-Distance-of-Profitability>) For 100MW, that would be \$300 million, which given the \$37million in annual fuel costs savings (electric vs gas costs for a year of 100MW), the system would pay for itself in less than 10 years. Regarding pollution produced, one can use the manufacturers data. For example the specs for the Purecell 400 (http://www.doosanfuelcell.com/attach_files/link/PureCell%20Model%20400%20Datasheet.pdf) or DFC3000 (http://www.fuelcellenergy.com/assets/PID000156_FCE_DFC3000_r3_hires.pdf). For the DFC3000 they are: NOx 0.01 lb/MWh SOx 0.0001 lb/MWh PM10 0.00002 lb/MWh For the Purecell 400 they are: NOx 0.01 lbs/MWh (0.006 kg/MWh) CO 0.02 lbs/MWh (0.009 kg/MWh) VOC 0.02 lbs/MWh (0.009 kg/MWh) SO2 Negligible Particulate Matter Negligible Water usage and discharge is significantly different. I would ask whether any experts in these technologies have been consulted and why this alternative has not been put in the application for consideration? I think this also points to a need to extend the deadline for this application since alternative technologies have to be evaluated carefully. This application should be rejected for it's incompleteness and limited solutions. What are the environmental impacts of the 100MW as DOM is

proposing? What is fueling the 100MW DOM is proposing to transmit to the facility? Where in this state/elsewhere is this pollution being generated and how does this compare to the efficiency and pollution of the fuel cells? While I am not sure if this is a good idea, I think it is in the best interests of all for the SCC to have all information as practical to make informed decisions. Andrew

160550042



160550042

Date: May 4, 2016

From: Edit Braunstein- Resident at Somerset Crossing Subdivision in Gainesville, VA

To: SCC

Subject: PUE-2015-00107. Haymarket 230 kV Double Circuit Transmission Line and 230-34.5 kV Haymarket Substation

SCC-CLERK'S OFFICE
STATE CORPORATION CONTROL CENTER:
2016 MAY 19 A 9:01

Mr. Chairman of the SCC,

My name is Edit Braunstein and, with my husband, we live at the Somerset Crossing neighborhood in Gainesville.

Back in 2007 we were looking for a house to buy when we moved from Florida State. We visited a beautiful, very affordable new house, being built at the moment and with opportunities to finalize it according to our desire in a Linton Hall subdivision; however when we looked at the surrounding area, very close to that house we saw power lines. From that moment that particular nice house was never again considered as a potential home for us. We finally settled for an older, more expensive house, in a quiet neighborhood, without the issues of power lines.

Nine years later we are coming not by our choice into a situation where the decision of having power lines close to our house is out of our control.

Fortunately, someone else other than two big corporations, Amazon and Dominion, have the power to stop this situation and it is you as a chairman of the SCC.

I hope and pray that you will have the clarity of mind to decide for what is the common sense solution and the common good for all the population affected by this situation: the Hybrid 66 proposal.

You see, all the other proposals will affect families and specially children living in the area of the project and beyond, with potential health issues.

The proposed route and the railroad route pass very close many homes and the two other alternative routes the Carver alternative route and the Madison alternative routes not only are close to many houses but are very close to The Elementary School. Children in the development age will be exposed to the EMF with potentially terrible consequences.

Even though there is not uniform agreement in the scientific community about the correlation between cancer and exposure to EMF, the danger cannot be dismissed.

In fact, the Australian Leukemia Foundation (2007) supports a prudent avoidance approach in relation to high-level EMF exposure (paper included) and also the British Journal of Cancer in an article published in 2013: Childhood leukemia close to high-voltage power lines--the Geocap Study, (108, pp. 1899-1906): we can read "In

conclusion, the present study has generated additional findings, based on a recent nationwide unselected population-based study, that support the hypothesis that living < 50m from a 225 / 400 kV HVOL may be associated with an increased incidence of childhood AL”.

We do not know what the new tools of biology and science in general will demonstrate in the future about these correlations, but we know that by choosing I-66 Hybrid proposal we will put behind a lot of concerns for all the families involved in this unwanted discussion and it could also avoid a lot of pain and suffering in the future.

Thanks for your attention.
Sincerely,

Edit Braunstein
Residence: 10507 Neale Sound Ct, Gainesville, VA 20155.

Biochemical Engineer and Supervisor of Manufacturing Science and Technology for
Cell Biology
ATCC- American Type Tissue Collection
10801 University Boulevard
Manassas, VA, 20110

Leukaemia Foundation of Australia



Position Statement: Powerlines and Childhood Leukaemia

Summary

The Leukaemia Foundation acknowledges that there is a great deal of community concern regarding the potential health effects of living close to high voltage powerlines. The possible link between proximity to these lines and exposure to the powerful electromagnetic fields (EMFs) they produce and childhood leukaemia has been the subject of a number of international studies over the past 30 years. While some have reported an *association* between exposure to high-level EMFs (greater than 0.4 μ T) and acute lymphoblastic leukaemia (ALL) in children, to date there is insufficient evidence to prove any *causal* link (i.e. that EMFs cause childhood leukaemia). There is no consistent evidence either to prove childhood exposures are associated with brain or any other types of solid tumours.

The scientific evidence surrounding the potential health effects of EMFs and in particular, magnetic fields remains uncertain. It cannot however be dismissed. As such the Leukaemia Foundation supports a *prudent avoidance* ⁶ approach in relation to high-level EMF exposure (see below).

Powerlines are part of the national grid that supplies electricity across Australia. They include high voltage transmission lines, which are mounted on large steel towers and lower voltage distribution lines mounted on concrete or wooden poles on the roadside, or placed underground.

Electromagnetic fields are a combination of electric and magnetic forces that travel together in wave-like patterns, at the speed of light. These waves travel at different frequencies. The frequency is simply the measurement of the number of times the electric and magnetic waves oscillate per unit of time. This is usually expressed in measurements known as Hz.

Electric fields result from electrical charges. The strength of the electric field is proportional to the voltage, i.e. the higher the voltage, the stronger the electric field. An electric field is produced as soon as an appliance is plugged in, regardless of whether it is switched on or not. Commonly used building materials such as wood and metal can be used to successfully shield us from these fields.

Magnetic fields result from the movement of electrical charges, also known as the current. The strength of the magnetic field is proportional to the current, i.e. the greater the amount of electricity flowing through a wire the stronger the magnetic field. Magnetic fields are commonly measured in units called tesla (T) or microtesla (μ T). A magnetic field is only generated when a device is switched on, or when an electric current flows through a wire. Magnetic fields pass easily through most types of material, making attempts at shielding both difficult and expensive.

Background

Electromagnetic fields (EMFs) are produced by all electrical equipment including electrical supply equipment (high and low voltage power lines, substations, above and below ground distribution cables, electrical wiring in business and homes), domestic appliances, TV and radiotransmitters, mobile phones and their base stations. They are also produced by natural forces in the earth. As such, it would be impossible for any of us to completely avoid exposure in today's world.

Exposure levels to EMFs from electric appliances around the home are generally in the range of 0.01 - 0.25 μ T. For homes near high voltage powerlines however, these levels may be as high as 0.5 - 1 μ T. For homes immediately under the high voltage powerline, magnetic field levels of 6 - 10 μ T may be found¹. It is important to point out that very few children in Australia live in close proximity to high-voltage powerlines. (the high-voltage distinction is important, as everyone has a powerline coming into their house)

There is a general consensus among experts, including the Australian radiation health authorities, that there is no scientific evidence to prove that ongoing exposure to electric and magnetic fields at the levels commonly found in the Australian environment (including most homes and work places), constitutes a *proven* health risk¹.

Both electric and magnetic fields are generally strongest close to their origin (for example a high-voltage power line) and become progressively weaker with distance. While the potential negative health effects of electric fields are generally thought to be limited to accidental electrocution, the potential health effects of magnetic fields are generally less well understood.

A comprehensive overview of EMFs and health is available at http://www.arpansa.gov.au/is_emf.htm

What does the research tell us?

The link between magnetic fields and childhood leukaemia has been investigated in a number of studies. In the most extensive and detailed to date, the UK Childhood Cancer Study (UKCCS)² found 'no evidence that either proximity to electrical installations or the magnetic field levels they produce in the UK is associated with increased risk of childhood leukaemia or any other cancer.' The researchers in this study measured children's residential exposure to electromagnetic fields (EMF) from a number of sources including domestic appliances, electrical wiring in homes, proximity to both low and high voltage power lines, underground power lines and electrical substations. They concluded that there was no evidence that exposure increased the risk of leukaemia, or any other type of childhood cancer.

In contrast, a recent UK study found that children who lived close to high voltage power lines at the time of their birth (within 200 m) had a significantly higher (1.69 times higher) chance of developing leukaemia than those who lived further away (beyond 600 m)³. Those who lived between 200-600 m of a line also had an increased risk (1.23 times higher) than those living further away. It is important to point out that the authors admitted to having 'no satisfactory explanation' for their findings. Exposure to magnetic fields from power lines was suggested as a *possible* explanation for the increased risk, however the authors pointed out that the strength of these forces was estimated to be typically less than the average magnetic fields found in most homes generated from other sources, such as domestic electrical appliances. The authors were also careful to state that the study does not *prove* that power lines cause leukaemia in children, and that there is no 'laboratory data or acceptable biological mechanism' to support their findings, which may have resulted from some other unexplained factors.

In a major European study⁴, researchers analysed the results from several previous studies of residential exposure to EMFs and childhood leukaemia. They found that a very small number of children (less than 1%) who were exposed to higher than average intensity magnetic fields (greater than 0.4 μ T) had around a two fold increased risk of childhood leukaemia, compared to those exposed to fields of less than 0.1 μ T. It is important to point out that very few children were found to be in this high-risk category, since very few children are ever exposed to such high intensity fields. Again the authors were unable to provide any 'explanation for the elevated risk' and suggest that at least some of it may have been due to internal problems with the study. Their findings however were consistent with an American study⁵, which identified a similarly increased risk for children exposed to fields 0.3 μ T or higher.

Prudent avoidance

Since 1991 the Energy Supply Association of Australia (ESAA), the peak organisation for electrical companies in Australia, has adopted a policy of *prudent avoidance* in regard to high-level EMF exposure. This involves taking exposure into consideration in the design and siting of new electrical facilities and avoiding where possible, and at a 'moderate cost and minimum inconvenience' homes, schools, playgrounds and 'other locations frequented by children'. It also involves sharing information with the public and involving them in decisions made regarding the siting of such facilities.

Some individuals may choose to take some simple steps to reduce their EMF exposure on a daily basis. These could include:

- Avoiding spending any time near high voltage powerlines or mobile phone base towers

- Switching off and unplug electrical equipment when not in use.
- Avoiding heated waterbeds.
- Avoiding electric blankets in beds, or at least switch them off and unplug before sleeping.
- Avoiding sleeping close to clock radios or any other electrical equipment.
- Sitting as far away as possible from your television or computer monitor.

If you are concerned about the level of EMF exposure in your home or workplace you can measure it using a special device available for hire from ARPANSA. For more information visit ARPANSA at www.arpansa.gov.au

Like many cancers, childhood ALL is thought to result from a series of changes in special proteins called genes, which normally control the growth and division of blood cells. The cause of changes remains unclear but there are likely to be a number of factors involved. For more information read the Leukaemia Foundation's *Understanding acute lymphoblastic leukaemia (ALL) in children - A guide for parents and families*.

References

Australian Radiation Protection and Nuclear Safety Agency 2003 The Controversy Over Electromagnetic Fields and Possible Adverse Health Effects. Fact sheet 8. Available at: http://www.arpansa.gov.au/is_emf.htm

UK Childhood Cancer Study Investigators. 2000 Childhood cancer and residential proximity to power lines. *Br J Cancer*, 83, 1573-80.

Draper GJ, Vincent T, Kroll ME and Swanson J. 2005 Childhood cancer in relation to distance from high-voltage power lines in England and Wales: a case-control study. *BMJ* Available from <http://www.bmj.com>.

Ahlbom A, Day N, Feychting M, et al. 2000 A pooled analysis of magnetic fields and childhood leukaemia. *Br J Cancer*, 83, 692-8.

Greenland S, Sheppard AR, Kaune WT, Poole C, Kelsh MA. A pooled analysis of magnetic fields, wire codes, and childhood leukemia. *Epidemiology* 2000;11: 624-34

World Health Organization 2000 Electromagnetic fields and public health cautionary policies. Available at: www.who.int/docstore/peh-emf/publications/facts_press/EMF-Precaution.htm

NRPB. Review of the Scientific Evidence for Limiting Exposure to Electromagnetic Fields (0- 300 GHz). *Doc NRPB*, 15, No. 3 (2004). Available at: www.hpa.org.uk/radiation/publications/documents_of_nrp/pdfs/doc_15_3.pdf

Keywords: childhood leukaemia; epidemiology; incidence; case-control; high-voltage overhead power lines

Childhood leukaemia close to high-voltage power lines – the Geocap study, 2002–2007

C Sermage-Faure^{1,2}, C Demoury^{1,2}, J Rudant^{1,2,3}, S Goujon-Bellec^{1,2,3}, A Guyot-Goubin^{1,2,3}, F Deschamps⁴, D Hemon^{1,2} and J Clavel^{*,1,2,3}

¹Inserm, Center for Research in Epidemiology and Population Health (CESP), U1018, Environmental Epidemiology of Cancer Team, Villejuif F-94807, France; ²Univ Paris-Sud, UMRS 1018, Villejuif F-94807, France; ³French National Registry of Childhood Hematological Malignancies (NRCH), Villejuif, France and ⁴Réseau de Transport d'Electricité (RTE), Coeur Défense, 110 esplanade du Général de Gaulle, La Defense 92030, France

Background: High-voltage overhead power lines (HVOLs) are a source of extremely low-frequency magnetic fields (ELF-MFs), which are classified as possible risk factors for childhood acute leukaemia (AL). The study was carried out to test the hypothesis of an increased AL incidence in children living close to HVOL of 225–400 kV (VHV-HVOL) and 63–150 kV (HV-HVOL).

Methods: The nationwide Geocap study included all the 2779 cases of childhood AL diagnosed in France over 2002–2007 and 30 000 contemporaneous population controls. The addresses at the time of inclusion were geocoded and precisely located around the whole HVOL network.

Results: Increased odds ratios (ORs) were observed for AL occurrence and living within 50 m of a VHV-HVOL (OR = 1.7 (0.9–3.6)). In contrast, there was no association with living beyond that distance from a VHV-HVOL or within 50 m of a HV-HVOL.

Conclusion: The present study, free from any participation bias, supports the previous international findings of an increase in AL incidence close to VHV-HVOL. In order to investigate for a potential role of ELF-MF in the results, ELF-MF at the residences close to HVOL are to be estimated, using models based on the annual current loads and local characteristics of the lines.

High-voltage overhead power lines (HVOLs) are one of the major sources of extremely low-frequency magnetic fields (ELF-MFs), considered a possible risk factor for childhood leukaemia. In the absence of any underlying biological hypothesis, the International Agency for Research on Cancer (IARC) classified ELF-MF as possible carcinogens (group 2B), based on epidemiological observations over more than two decades (IARC, 2002). The first meta-analyses concluded that exposure to ELF-MF levels of at least $0.3 \mu\text{T}$ was significantly associated with an increased incidence of childhood acute leukaemia (AL) (odds ratio (OR) = 1.7 (1.2–2.3) for exposures $\geq 0.3 \mu\text{T}$ (Greenland *et al*, 2000) and OR = 2.0 (1.3–3.3) for exposures $\geq 0.4 \mu\text{T}$ (Ahlbom *et al*, 2000)). A recent meta-analysis of the studies published after 2000 (Kheifets *et al*, 2010) generated consistent but weaker results (OR = 1.4 (0.9–2.4) for exposures $\geq 0.3 \mu\text{T}$). The large British study by Draper *et al* (2005) focused on the proximity of VHV-HVOL and showed an

association between AL and residence at birth <200 m from a VHV-HVOL (OR = 1.7 (1.1–2.5)) and, to a lesser extent, between 200 and 600 m from a VHV-HVOL (OR = 1.2 (1.0–1.5)). With the same data, the relative risk was not significantly increased for estimates of ELF-MF $\geq 0.4 \mu\text{T}$ (Kroll *et al*, 2010). High-voltage overhead power lines account for only a fraction of ELF-MF exposure, but, in their near vicinity, constitute the main source of background exposure (Schüz *et al*, 2000; Maslanyj *et al*, 2007).

The aim of the present study was to test whether the risk of AL was increased in the vicinity of HVOL, where children were expected to encounter higher residential exposure to ELF-MF. We followed a two-step approach. The present one aims at investigating the relationship between AL and distance to HVOL. The second step will rely on calculated residential exposure to ELF-EMF based on characteristics of the neighbouring HVOL. The study, the first in France, was based on the geolocation of the last

*Correspondence: Dr J Clavel; E-mail: Jacqueline.Clavel@inserm.fr

Received 13 November 2012; revised 25 February 2013; accepted 2 March 2013; published online 4 April 2013

© 2013 Cancer Research UK. All rights reserved 0007–0920/13

address and covered the entire mainland over a recent period (2002–2007), on an exhaustive basis, free from participation bias, and was based on a geographic information system (GIS) using precise and recent databases to locate the dwellings and HVOL.

MATERIALS AND METHODS

The Geocap case-control study. The Geocap case-control study included all the 2779 French childhood AL cases aged < 15 years at the end of the year of diagnosis, diagnosed between 1 January 2002 and 31 December 2007, residing in mainland France (excluding Corsica for which HVOL information was not available). The cases were obtained from the French National Registry of Childhood Hematopoietic Malignancies (Lacour *et al*, 2010).

Over the same period (2002–2007), six yearly sets of 5000 control addresses were randomly sampled from the paediatric population of mainland France by the National Institute for Statistics and Economic Studies (INSEE), using the income and council tax databases of the French households. These databases contain the addresses and income information of all the households in France, irrespective of employment status, and list the children in each household by year of birth. The sample was stratified on the 94 French mainland administrative areas (*Départements*). The individual variables available for the controls were the year of birth, number of children in the household and last address. Demographic and socioeconomic characteristics of the municipality (*Commune*) of residence were also used as contextual variables. The sample of 30 000 controls was closely representative of the source population in terms of age and number of children in the household, and in terms of contextual variables, that is, size of the urban unit, median income, proportion of blue-collar workers, proportion of subjects who successfully completed high school (baccalaureate holders) and proportion of homeowners in the *Commune* of residence (Sermage-Faure *et al*, 2012).

Geocoding. The residence considered for geolocation was the residence at the time of diagnosis for the cases and that at the time of inclusion for the controls. Residential histories, particularly addresses at birth, were not available. The method for geocoding the addresses of cases and controls was compiled, checked for consistency and corrected when necessary by GEOCIBLE, an outside service provider, in close cooperation with the epidemiology research team. The addresses were geocoded blind to case/control status, using the MAPINFO GIS, NAVTEQ street databases and detailed vectorized maps from the National Geographic Institute (IGN). Automatic processes were checked and completed by visual inspection of maps when necessary. Ultimately, only 3% of the cases and 1% of the controls could not be located more precisely than by their *Commune* of residence and were allocated the coordinates of their *Commune* town hall.

In the Navteq and IGN databases, the geocodes are given at the middle of the street in front of the number in the street (i.e., the front door, the entrance of the plot or the projection of centre of the plot along the street), generally corresponding to the mailbox residence. Most often in urban areas and in collective housing, the mailbox is attached to the building of residence. However, especially in countryside, the house can be at a distance from the entrance of the plot, where the mailbox is.

Depending on whether the databases enabled location of the home directly or by extrapolation from the nearest or more distant neighbours, the coordinates were assigned a degree of uncertainty along the street ranging from 20 m (exact number in the database) to the size of a *Commune* (Table 1). The scale of uncertainty provided by Geocible had been determined previously, based on the size of the objects to locate and on the mean differences between estimated and measured geocodes. The best geocoded

Table 1. Distribution of the cases and controls by category of uncertainty of location by geocoding

Category of accuracy of address location for geocoding	Uncertainty	Cases		Controls	
		N	%	N	%
At the exact number	20 m	1946	70.0	23 171	77.2
In a section of a short street	50 m	173	6.2	1658	5.5
At a close number	100 m	130	4.7	801	2.7
In a medium street or in a hamlet	300 m	394	14.2	3693	12.3
In a long street	500 m	54	1.9	383	1.3
In a <i>Commune</i>		82	3.0	292	1.0
Total		2779	100.0	30 000	100.0

addresses were assigned an uncertainty of 20 m, equal to the mean value of the estimated coordinates given by the GIS and the center of the house. Altogether, 1946 cases (70%) and 23 171 controls (77%) were located by their exact number in the street (best geocoded addresses, uncertainty of 20 m), whereas 303 cases (11%) and 2459 controls (8.6%) were located by a segment of a short street or by a close number (uncertainty of 50–100 m).

In addition to the coordinates obtained for all the postal addresses of the Geocap sample, another set of coordinates was also estimated using photographic views obtained from Street View (Google Maps), Géoportail (IGN data) and the French cadaster, when available. This was possible for 72% of the cases and 69% of the controls living < 200 m from a HVOL, considering the uncertainty, and used to position the building of residence.

HVOL characteristics and distance from the nearest HVOL. There are 77 400 km of HVOL in France. There are five main types: HVOL of 400 kV (13 350 km), 225 kV (21 200 km), 150 kV (1050 km) and 90 or 63 kV (41 800 km). The HVOLs have been precisely mapped by RTE (*Réseau de Transport d'Electricité*), the French utility in charge of electricity transmission, based on the utility's own database, in which pylons and sections of lines are accurately located, and on the most precise local maps of the national geographic institute (IGN). The distances from the closest HVOLs were estimated by GEOCIBLE.

Statistical analysis. All the statistical analyses were performed using the SAS software package (version 9; SAS Institute Inc., Cary, NC, USA). The ORs, their 95% confidence intervals and two-sided *P*-values were estimated by unconditional logistic regression adjusted for age in 5-year categories and *Département*. Additional analyses stratified by age category were adjusted for age in years.

The subjects were classified in terms of their distance from the closest HVOL (< 50, 50–99, 100–199, 200–599 and ≥ 600 m). The very high voltage lines, 225 and 400 kV (VHV-HVOL), and the high voltage lines, 63, 90 and 150 kV (HV-HVOL), were separated as dwellings located ≤ 50 m from VHV-HVOL are expected to be more frequently exposed to higher ELF-MF than those located ≤ 50 m from HV-HVOL (Maslanyj *et al*, 2009). The inverse distance function was used to test for the existence of a trend in AL incidence, assigning 0 to the dwellings located at least 600 m from an HVOL.

All the main analyses were conducted on the whole study sample, without any selection by address uncertainty. The analyses

Table 2. Description of the cases included in the study, by the distance between their residence and the closest HVOL and by voltage category (very high (225–400 kV) or high (63–150 kV)) over the period 2002–2007

	Distance from the closest HVOL						Total
	225–400 kV HVOL			63–150 kV HVOL			
	0–49 m	50–199 m	≥200 m	0–49 m	50–199 m	≥200 m	
Gender							
Female	4	12	1246	7	26	1229	1262
Male	5	12	1500	7	28	1482	1517
Age							
<5 years	6	11	1291	8	24	1276	1308
5–9 years	3	10	858	2	18	851	871
10–14 years	0	3	597	4	12	584	600
Down's syndrome	0	0	43	0	0	43	43
AL type							
ALL	8	21	2250	13	46	2220	2279
B-cell precursor ALL	6	12	1056	8	25	1041	1074
T-cell ALL	0	0	173	1	4	168	173
Other ALL	2	9	1021	4	17	1011	1032
AML	1	2	428	1	6	424	431
Other AL	0	1	68	0	2	67	69

Abbreviations: AL = acute leukaemia; ALL = acute lymphoblastic leukaemia; AML = acute myeloblastic leukaemia; HVOL = high-voltage overhead power line.

were conducted on all the cases and also stratified by age group – <5 years old covering most of the incidence peak and ≥5 years old – and for acute lymphoblastic leukaemia (ALL) alone.

All the analyses were performed taking the same baseline as reference category, that is, the group of children who lived in *Communes* with no part of their territory within 600 m of a HVOL, after accounting for geocoding uncertainty. Thus, the baseline included all the residences definitely located ≥600 m from a HVOL, even if they were geocoded with the highest uncertainty. Additional sensitivity analyses also included the subjects living at least 600 m from a HVOL in the reference category, even when the *Commune* had a part of its territory within 600 m of a HVOL, in order to account for the possibility that the baseline category might select residences in relation with an AL risk factor.

The 67 cases and 203 controls who lived in a *Commune* partially located within 600 m of a line but who could not be individually located better than at the town hall were considered to have missing data for the distances from HVOL.

Supplementary analyses were performed to test the robustness of the results and account for the spatial extent of the house, by restriction to the best geocoded addresses (uncertainty <20 m), or by modifying the distance cutoffs around the *a priori* value of 50 m (30, 40, 60 and 70 m). In addition, for sensitivity analyses, when the distances using the main geocoding and the photographic views were available, the cases and controls were classified in the category '<50 m from a HVOL', either when the distance from photographic views was <50 m, or when at least one of the two estimated distances was <50 m, or when both the estimated distances were <50 m.

The analyses were also stratified by contextual socioeconomic variables extracted from the 1999 census data for the *Commune* of residence, including the urban status of the *Commune*, median income of the households, proportion of blue-collar workers and proportion of baccalaureate holders. Additional analyses were performed after excluding the cases and controls who lived <5 km

from a nuclear power plant, in order to rule out possible confounding by residence in the proximity of a nuclear power plant, which was associated with AL in the present study (Sermage-Faure *et al*, 2012).

RESULTS

Table 2 describes the cases registered from 2002 to 2007 by age, gender and leukaemia subtype, on the basis of the distance of their residences from the closest VHV-HVOL or HV-HVOL.

The 610 cases (22.0%) and 7061 controls (23.5%) who were living in a *Commune* entirely located at least 600 m from any HVOL constituted the baseline of the models. Living within 50 m of the closest HVOL, all voltages considered together, was not associated with AL (OR = 1.2 (0.8–1.9)) (Table 3). However, while no association was observed with residences close to HV-HVOL (OR = 1.0 (0.6–1.7)), an association was evidenced for children who lived within 50 m of a VHV-HVOL (OR = 1.7 (0.9–3.6)). In contrast, the ORs were close to one for the residences located ≥50 m from a HVOL, even a VHV-HVOL, and no statistically significant trend was observed with the inverse of the distance ($P = 0.28$ for distance from VHV-HVOL). The results for ALL were very similar (OR = 1.9 (0.9–4.0) at <50 m from a VHV-HVOL).

Splitting the sample into children aged <5 years and those aged ≥5 years showed that the association was only observed for the younger group (Table 4). In that age group, living within 50 m of the closest VHV-HVOL was significantly associated with AL (OR = 2.6 (1.0–7.0)), with a significant trend with the inverse of the distance ($P = 0.03$), whereas there was no association for the older group (OR = 1.0 (0.3–3.3) living within 50 m of the closest VHV-HVOL).

Living within 50 m of a VHV-HVOL was not associated with AL in the *Communes* of urban units with a population >100 000

Table 3. Association between childhood acute leukaemia and distance to the closest HVOL by category of voltage (very high (225–400 kV) or high (63–150 kV)) over the period 2002–2007

	Distance to HVOL																	
	225–400 kV HVOL						63–150 kV HVOL						Any HVOL					
	Cases		Controls		OR ^a	95% CI	Cases		Controls		OR ^a	95% CI	Cases		Controls		OR ^a	95% CI
n	%	n	%	n			%	n	%	n			%	n	%	n		
Baseline ^b	610	22.0	7061	23.5	1.0	Reference	610	22.0	7061	23.5	1.0	Reference	610	22.0	7061	23.5	1.0	Reference
Unknown	67		203				67		203				67		203			
≥600 m	1924	69.2	20896	69.7	1.0	(0.9–1.2)	1792	64.5	19168	63.9	1.1	(1.0–1.2)	1665	59.9	17937	59.8	1.1	(0.9–1.2)
200–599 m	145	5.2	1416	4.7	1.2	(1.0–1.4)	242	8.7	2740	9.1	1.0	(0.8–1.2)	345	12.4	3633	12.1	1.1	(0.9–1.2)
100–199 m	16	0.6	267	0.9	0.7	(0.4–1.2)	33	1.2	461	1.5	0.8	(0.6–1.2)	44	1.6	669	2.2	0.8	(0.5–1.0)
50–99 m	8	0.3	97	0.3	1.0	(0.5–2.1)	21	0.8	203	0.7	1.2	(0.7–1.9)	25	0.9	284	0.9	1.0	(0.7–1.6)
0–49 m	9	0.3	60	0.2	1.7	(0.9–3.6)	14	0.5	164	0.5	1.0	(0.6–1.7)	23	0.8	213	0.7	1.2	(0.8–1.9)
Total	2779		30000				2779		30000				2779		30000			

Abbreviations: CI = confidence interval, HVOL = high-voltage overhead power line; OR = odds ratio.

^aOR and 95% CIs estimated by logistic regression adjusted for age at the end of the year (5-year age groups for the 0–14-year-old children, 1-year age groups for the 0–4-year-old children) and Département of residence.

^bResidence in a Commune entirely located ≥600 m from an HVOL.

(Table 4), but an association was observed for the less urban categories of *Commune*. The same pattern was observed for the under-5-year age group (data not shown). The association between AL and living <50 m from a VHV-HVOL appeared more marked, although not significantly so, in the *Communes* with less-favorable contextual socioeconomic characteristics: median income or percentage baccalaureate holders lower than the median value for the controls; percentage blue-collar workers greater than the median value for the controls. Adjustments for those contextual variables, either separately or jointly, did not change the estimates.

No case and only two controls lived within 5 km of a nuclear power plant and <200 m from a VHV-HVOL; excluding them did not modify the results.

Sensitivity analyses restricted to the best geocoded subjects (uncertainty ≤20 m) generated slightly stronger results (OR = 2.1 (0.9–4.7) for living within 50 m of a VHV-HVOL) (Table 5). The results were also unchanged when the cutoffs were 10 and 20 m above or below the *a priori* value of 50 m, and when the baseline was extended to include the subjects living >600 m from a HVOL, even if their *Commune* of residence had parts located <600 m from a line (data not shown). Lastly, in the sensitivity analyses using the main geocoding distance and that based on photographic views when both were available (Table 5), the ORs remained of the same order of magnitude but the associations were no longer significant (OR = 1.3 (0.5–3.7) for distance <50 m based on photographic views, OR = 1.7 (0.6–4.8) for both distances <50 m and OR = 1.5 (0.8–3.1) for at least one distance <50 m from a VHV-HVOL). For 0–4-year-old children, this was also the case (OR = 2.5 (0.6–10.5) for distance <50 m based on photographic views, OR = 3.5 (0.8–15.1) for both distances <50 m and OR = 2.3 (0.9–6.0) for at least one distance <50 m).

DISCUSSION

The present analysis of the Geocap nationwide case-control study was carried out to test the hypothesis that living close to HVOL, particularly VHV-HVOL, was associated with an increased incidence of childhood AL. The study focused on HVOL, a

major source of exposure to ELF-MF in neighbouring residences (Schüz *et al*, 2000; Maslanyj *et al*, 2007). The proximity of HVOL to the residence of all the subjects was reliably evaluated without any selection and using the same process over all mainland France and over the 2002–2007 period. The results for living <50 m from a 225 or 400 kV HVOL were compatible with the IARC conclusions. There was no association beyond that distance. The association at a short distance was not observed for children aged ≥5 years or those living in the most urban *Communes*.

The study covered a recent and relatively short period, and historical databases were therefore available for the entire period. One of the main strengths of the Geocap study is that it was designed to avoid selection biases. The cases were identified by the national registry, which complies with the international criteria required for cancer registration and classification, and achieves a high degree of completeness, by active research with almost three sources per case on average (Clavel *et al*, 2004; Lacour *et al*, 2010). Similarly, the recruitment of the controls did not require their active participation, preventing self-selection by socioeconomic status. *De facto*, the control sample was closely representative of the paediatric population on the basis of the sociodemographic contextual criteria (Table 1).

All the cases' and controls' addresses were obtained and geocoded, and the distances were calculated from objective databases free from any recall bias and blind to case/control status. This is another strength of the Geocap study in that it enabled minimisation of differential misclassifications. The distances estimated from the GISs were assumed to rank, as adequately as possible, the cases and controls by the true distance of their dwellings from the HVOL. The databases used to locate the lines were very precise. In particular, pylons were located with an uncertainty of 2.5 m in the RTE database.

Interestingly, the results were strengthened when the analyses were restricted to the best geocoded addresses. The 67 cases and 203 controls (<2% of the subjects) whose addresses were not precise enough to enable their location close to the HVOL probably had no substantial impact on the results, given the expected distribution of the few subjects with respect to the distance from VHV-HVOL (about 0.2% of the controls <50 m from VHV-HVOL). For the association to have been due to the

Table 4. Association between childhood AL and distance from HVOLs over the period 2002–2007, stratified by age and urban status of the *Commune* of residence

	Address-based distance from HVOL											
	225–400 kV HVOL						63–150 kV HVOL					
	Ca		Co		OR ^a	95% CI	Ca		Co		OR ^a	95% CI
	n	%	n	%			n	%	n	%		
Age												
0–4 years												
Baseline ^b	311	23.8	2326	23.9	1.0	Reference	311	23.8	2326	23.9	1.0	Reference
Unknown	35		85				35		85			
≥600 m	870	66.6	6734	69.3	0.9	(0.8–1.1)	814	62.3	6146	63.2	0.9	(0.8–1.1)
200–599 m	74	5.7	444	4.6	1.2	(0.9–1.6)	115	8.8	902	9.3	0.9	(0.7–1.1)
100–199 m	5	0.4	87	0.9	0.4	(0.2–1.0)	13	1.0	145	1.5	0.6	(0.4–1.2)
50–99 m	6	0.5	27	0.3	1.6	(0.7–4.1)	11	0.8	61	0.6	1.3	(0.7–2.5)
0–49 m	6	0.5	14	0.1	2.6	(1.0–6.9)	8	0.6	52	0.5	1.1	(0.5–2.3)
Total	1307		9717				1307		9717			
5–14 years												
Baseline ^b	299	20.3	4735	23.3	1.0	Reference	299	20.8	4735	23.3	1.0	Reference
Unknown	32		118				32		118			
≥600 m	1054	71.6	14 162	69.8	1.2	(1.0–1.4)	978	66.4	13 022	64.2	1.2	(1.0–1.4)
200–599 m	71	4.8	972	4.8	1.2	(0.9–1.5)	127	8.6	1838	9.1	1.1	(0.9–1.4)
100–199 m	11	0.7	180	0.9	1.0	(0.5–1.8)	20	1.4	316	1.6	1.0	(0.6–1.6)
50–99 m	2	0.1	70	0.3	0.5	(0.1–2.0)	10	0.7	142	0.7	1.1	(0.6–2.1)
0–49 m	3	0.2	46	0.2	1.0	(0.3–3.3)	6	0.4	112	0.6	0.9	(0.4–2.1)
Total	1472		20 283				1472		20 283			
Size of urban unit^c												
<5000 inhabitants												
Baseline ^b	309	34.1	3415	35.5	1.0	Reference	309	34.1	3415	35.5	1.0	Reference
Unknown	36		67				36		67			
≥600 m	525	57.9	5630	58.6	1.0	(0.9–1.2)	482	53.1	5221	54.3	1.0	(0.9–1.2)
200–599 m	30	3.3	393	4.1	0.9	(0.6–1.3)	68	7.5	724	7.5	1.0	(0.8–1.4)
100–199 m	2	0.2	71	0.7	0.3	(0.1–1.4)	7	0.8	109	1.1	0.7	(0.3–1.6)
50–99 m	1	0.1	20	0.2	0.6	(0.1–4.6)	4	0.4	45	0.5	1.1	(0.4–3.0)
0–49 m	4	0.4	19	0.2	2.5	(0.8–7.7)	1	0.1	34	0.4	0.4	(0.1–2.9)
Total	907		9615				907		9615			
5000–100 000 inhabitants												
Baseline ^b	71	11.1	756	10.9	1.0	Reference	71	11.1	756	10.9	1.0	Reference
Unknown	18		63				18		63			
≥600 m	513	80.3	5811	84.1	0.8	(0.6–1.1)	451	72.7	4942	71.6	0.9	(0.7–1.2)
200–599 m	27	4.2	219	3.2	1.2	(0.7–2.0)	72	11.5	876	12.7	0.8	(0.5–1.1)
100–199 m	5	0.8	33	0.5	1.5	(0.5–3.9)	15	2.4	144	2.1	1.0	(0.5–1.8)
50–99 m	1	0.2	19	0.3	0.6	(0.1–4.8)	5	0.8	73	1.1	0.7	(0.3–1.7)
0–49 m	4	0.6	6	0.1	4.9	(1.3–19.2)	7	1.1	53	0.8	1.2	(0.5–2.7)
Total	639		6907				639		6907			
≥100 000 inhabitants												
Baseline ^b	230	18.7	2890	21.4	1.0	Reference	230	18.7	2890	21.4	1.0	Reference
Unknown	13		73				13		73			
≥600 m	886	71.9	9455	70.2	1.1	(0.9–1.4)	859	69.7	9005	66.8	1.2	(0.9–1.4)
200–599 m	88	7.1	804	6.0	1.3	(1.0–1.8)	102	8.3	1140	8.5	1.1	(0.8–1.4)
100–199 m	9	0.7	163	1.2	0.7	(0.4–1.4)	11	0.9	208	1.5	0.7	(0.4–1.3)
50–99 m	6	0.5	58	0.4	1.4	(0.6–3.3)	12	1.0	85	0.6	1.7	(0.9–3.3)
0–49 m	1	0.1	35	0.3	0.4	(0.1–2.9)	6	0.5	77	0.6	1.0	(0.4–2.4)
Total	1233		13 478				1233		13 478			

Abbreviations: AL = acute leukaemia; Ca = number of cases; CI = confidence interval; Co = number of controls; HVOL = high-voltage overhead power line; OR = odds ratio.

^aORs and 95% CIs estimated by logistic regression adjusted for age at the end of the year (5-year age groups for the 0–14-year-old children, 1-year age groups for the 0–1-year-old children) and *Département* of residence.

^bBaseline = residence in a *Commune* entirely located ≥600 m from any HVOL.

^cAn urban unit is defined by the INSEE (National Institute of Statistics and Economic Studies) as a group of *Communes* in which the distance between dwellings is nowhere more than 200 m.

Table 5. Sensitivity analyses of the association between childhood acute leukaemia and distance to the closest HVOL by category of voltage (very high (225–400 kV) or high (63–150 kV)) over the period 2002–2007, for all ages and the 0–4-years age group

Main results				Sensitivity analyses																
GIS with any uncertainty				(1) GIS with uncertainty of 20 m				(2) Photographic views available												
								Photograph only				<50 m by GIS or photo				<50 m by GIS and photo				
Ca	Co	OR	95% CI	Ca	Co	OR	95% CI	Ca	Co	OR	95% CI	Ca	Co	OR	95% CI	Ca	Co	OR	95% CI	
0–14 years																				
Baseline	610	7061	1.0	Reference	610	7061	1.0	Reference	610	7061	1.0	Reference	610	7061	1.0	Reference	610	7061	1.0	Reference
Any HVOL																				
100–199 m	44	669	0.8	(0.5–1.0)	32	499	0.7	(0.5–1.1)	36	455	0.9	(0.6–1.3)								
50–99 m	25	284	1.0	(0.7–1.6)	20	212	1.1	(0.7–1.8)	22	184	1.4	(0.9–2.2)								
0–49 m	23	213	1.2	(0.8–1.9)	16	152	1.2	(0.7–2.0)	14	158	1.1	(0.6–1.9)	24	243	1.2	(0.7–1.8)	13	128	1.2	(0.7–2.1)
VHV-HVOL																				
100–199 m	16	267	0.7	(0.4–1.2)	13	200	0.8	(0.4–1.3)	18	172	1.2	(0.7–1.9)								
50–99 m	8	97	1.0	(0.5–2.1)	6	68	1.0	(0.4–2.4)	8	74	1.3	(0.6–2.7)								
0–49 m	9	60	1.7	(0.9–3.6)	7	39	2.1	(0.9–4.7)	4	38	1.3	(0.5–3.7)	9	68	1.5	(0.8–3.1)	4	30	1.7	(0.6–4.8)
HV-HVOL																				
100–199 m	33	461	0.8	(0.6–1.2)	22	346	0.7	(0.5–1.1)	24	322	0.9	(0.6–1.3)								
50–99 m	21	203	1.2	(0.7–1.9)	16	155	1.2	(0.7–2.0)	14	121	1.3	(0.7–2.3)								
0–49 m	14	164	1.0	(0.6–1.7)	9	120	0.8	(0.4–1.7)	10	127	0.9	(0.5–1.8)	15	188	0.9	(0.5–1.6)	9	103	1.0	(0.5–2.0)
0–4 years																				
Baseline	311	2326	1.0	Reference	311	2326	1.0	Reference	311	2326	1.0	Reference	311	2326	1.0	Reference	311	2326	1.0	Reference
Any HVOL																				
100–199 m	17	213	0.6	(0.3–1.0)	9	160	0.4	(0.2–0.8)	16	146	0.9	(0.5–1.5)								
50–99 m	14	84	1.2	(0.7–2.1)	11	62	1.2	(0.6–2.4)	13	55	1.7	(0.9–3.3)								
0–49 m	14	62	1.5	(0.8–2.8)	9	42	1.3	(0.6–2.8)	8	38	1.5	(0.6–3.3)	14	66	1.4	(0.8–2.6)	8	34	1.7	(0.8–3.9)
VHV-HVOL																				
100–199 m	5	87	0.4	(0.2–1.0)	4	63	0.5	(0.2–0.4)	9	58	1.3	(0.6–2.8)								
50–99 m	6	27	1.6	(0.7–4.1)	4	19	1.6	(0.5–5.0)	6	21	2.1	(0.8–5.8)								
0–49 m	6	14	2.6	(1.0–6.9)	5	8	4.1	(1.3–13.3)	3	7	2.5	(0.6–10.5)	6	16	2.3	(0.9–6.0)	3	5	3.5	(0.8–15.1)
HV-HVOL																				
100–199 m	13	108	0.6	(0.4–1.2)	6	145	0.4	(0.2–0.9)	10	99	0.7	(0.4–1.4)								
50–99 m	11	45	1.3	(0.7–2.5)	8	61	1.2	(0.5–2.6)	7	35	1.5	(0.6–3.4)								
0–49 m	8	36	1.1	(0.5–2.3)	4	52	0.6	(0.2–1.8)	5	33	1.1	(0.4–2.9)	8	55	1.0	(0.5–2.1)	5	30	1.3	(0.5–3.4)

Abbreviations: Ca = number of cases; CI = confidence interval; Co = number of controls; GIS = geographic information system; HV-HVOL = high voltage high-voltage overhead power lines (63–150 kV); HVOL = high-voltage overhead power line; OR = odds ratio; VHV-HVOL = very high voltage high-voltage overhead power lines (225–400 kV). The first sensitivity analysis (1) is restricted to the addresses best geocoded (GIS obtained with uncertainty of 20 m) and the second one (2) to the addresses for which a photographic view was available. The results shown in Tables 4 and 5 are recalled in the first columns.

Unknown category, the true addresses would have to have been within 50 m of a VHV-HVOL for none of the unclassified cases and for about 15% of the unclassified controls, which is very unlikely. The sensitivity analyses were consistent with the main results.

The Geocap study was designed to avoid selection and differential misclassification biases, which are common shortcomings of case-control studies on environmental factors, particularly ELF-MF (Mezei and Kheifets, 2006; Kheifets and Oksuzyan, 2008; Schüz and Ahlbom, 2008). The study included no individual data other than age and address, which were obtained for all the cases and controls. Therefore, potential AL risk factors such as birth order, breastfeeding, day-care attendance and pesticide exposure were not available. However, conditionally on age and the sociodemographic characteristics of the Commune of residence, which were accounted for in adjusted or stratified analyses, known or suspected risk factors are not likely to differ markedly within vs outside the 50-m distance from the VHV-HVOL. The study may have suffered from non-differential misclassifications, particularly because of the uncertainty of the geolocation of the homes, or because the period considered, that is, residence at diagnosis or interview, may not belong to the most relevant time window, or because the small numbers did not enable

separation of the 400- and 225-kV VHV-HVOL or splitting the smallest category of distance. Therefore, the relationship between living close to VHV-HVOL and AL is probably not overestimated. As a registry-based study, the Geocap study considered the addresses at the time of diagnosis for the cases and at the time of inclusion for the controls. It did not cover the whole residential history since conception, and earlier or longer time windows may be more relevant in childhood AL. In the Escale case-control study (data collected in 2003–2004), the household had not moved during the index pregnancy or childhood for 46% of the controls < 15 years, and 60% of those < 5 years (Amigou *et al*, 2011). In the present study, the relationship was only observed for children < 5 years, which might be compatible with a smaller impact of misclassifications, due to moves, of early exposures related to the proximity of VHV-HVOL. The relationship was not observed in children living in the most populated urban Communes.

The present study exclusively addressed the question, recurrent in France, of the risk of childhood AL close to HVOL. If living < 50 m from HVOL is causally related to AL, it is expected to induce an excess of less than one new case < 15 years per year in France, under steady conditions of residency close to VHV-HVOL. The distance of the residence from a HVOL is by no means a perfect surrogate for individual exposure to ELF-MF because of the

proximity of the lines. Individual *in situ* measurements would be more suitable exposure indicators, provided that they were standardized, accurate and precise measurements, and that no selection bias (and no participation bias) limited their interpretation. Residential proximity of a VHV-HVOL was considered an indicator of increased probability of high residential exposure to ELF-MF, with the hypothesis that other sources of exposure to ELF-MF would be independent of the presence of the line and thus would be distributed similarly for the children living <50 m from a VHV-HVOL and those living further away (Schüz *et al*, 2000; Maslanyj *et al*, 2007). The study combined stringent voltage (≥ 225 kV) and distance (<50 m) conditions with a high degree of accuracy in the geocoding process, in order to identify the individuals who most probably had the highest exposures to ELF-MF in the population study. Exposure to ELF-MF depends on many sources and, regarding power lines, on many other parameters than distance, particularly current load and type of pylon (also related to the line voltage). Conversely, the distance from VHV-HVOL might also be an indicator of environmental exposures and lifestyle factors related to the vicinity of lines other than ELF-MF.

In a descriptive analysis of studies of ELF-MF exposure in 4452 homes in the United Kingdom (UKCCS, 1999) and 1835 homes in Germany (Schüz *et al*, 2000), only a small number of dwellings were located within 50 m of a HVOL (93 homes), 16 of which were close to a 220–400 kV HVOL (Maslanyj *et al*, 2009). Extremely low-frequency magnetic field exposure $\geq 0.4 \mu\text{T}$ was more prevalent in the latter homes (18.8%) than in those close to 11–132 kV HVOL (6.5%), even though the absolute numbers of dwellings with ELF-MF exposure $\geq 0.4 \mu\text{T}$ were similar (three and five homes, respectively). Therefore, in this study, the absence of an association close to HV-HVOL lines, where the prevalence of exposed residences is assumed to be lower, is poorly informative with respect to the hypothesis that ELF-MF may have a role in childhood AL.

This hypothesis will be investigated more precisely in a future stage of the Geocap study. RTE is to calculate individual estimates of the exposure to ELF-MF for all the Geocap subjects located close to a HVOL, blind to case/control status. The exposure estimates will take into account the particular characteristics of each of the neighbouring lines (pylons geometry, height and type of cable, ground wires and so on), the average annual current load for each of the identified lines, the time-distribution percentiles of the current load and the particular location of the residence with respect to the closest line spans (Bessou *et al*, 2013).

This is the first French contribution to the issue of ELF-MF, HVOL and childhood AL. The results are compatible with the first meta-analyses published in 2000 (Ahlbom *et al*, 2000; Greenland *et al*, 2000), the recent review by Schüz and Ahlbom (2008) and the most recent meta-analysis summarizing the studies of the last decade (Kheifets *et al*, 2010). While no underlying biological mechanism has been advanced to date in support of the epidemiological observation (WHO, 2007), the IARC classification of ELF-MF as a possible carcinogen (IARC, 2002) has not been strongly challenged. The study by Draper *et al* (2005) based on residence at birth and covering more than three decades (1962–1995) revealed associations with longer distances from power lines than previously envisaged, far above the threshold usually recognised as generating ELF-MF greater than background exposures, and with a positive trend with decreasing distance. Extremely low-frequency magnetic fields were estimated secondarily in the same study, and then considered unlikely to be the only explanation (Swanson, 2008; Kroll *et al*, 2010) for the observed relationship with distance. Overall, the number of exposed newborns was small because five AL cases and three controls resided at birth within 50 m of a HVOL (mainly VHV-HVOL) (Draper *et al*, 2005), and two AL cases and one control were

assumed to be exposed to at least $0.4 \mu\text{T}$ (Swanson, 2008). In the present study, we observed no significant trend with decreasing distance to VHV-HVOL.

Recently, in a commentary on the most recent papers by Kroll *et al* (2010) and Kheifets *et al* (2010), Schmiedel and Blettner (2010) drew attention to the current limitations of epidemiology with regard to affording new insights in the field and answering questions in the absence of satisfactory biological models. Geocap was designed for quantitative modelling and the study of coexposures, and may thus be considered an appropriate tool for contributing to knowledge in the field.

CONCLUSION

In conclusion, the present study has generated additional findings, based on a recent nationwide unselected population-based study, that support the hypothesis that living <50 m from a 225 or 400 kV HVOL may be associated with an increased incidence of childhood AL. No increase in risk was observed further from those lines and no increase in childhood AL risk was detected within 50 m of the 63–150 kV HVOL. Model-based estimates of ELF-MF exposures will be used to investigate for potential involvement of ELF-MF in the observed association.

ACKNOWLEDGEMENTS

This study was supported by the Institut de Veille Sanitaire (InVS), the Agence Nationale de Sécurité Sanitaire de l'Alimentation, de l'Environnement et du Travail (ANSES), the Fondation ARC pour la recherche sur le cancer, the Fondation Pfizer, the Institut National du Cancer (INCa), the Agence Nationale de la Recherche (ANR) and the Cancéropôle Ile de France. It is part of the Investissement d'Avenir HOPE-EPI. The authors are particularly grateful to: (1) Olivier Lamy and Mathieu Carrère from 'GEOCIBLE', who carefully ensured all the geocoding, and IGN, which made available precise maps for the whole country; (2) Magda Tomasini and Laurent Auzet from the 'Institut National de la Statistique et des Etudes Economiques' (INSEE), who conducted the control sampling; (3) Jérôme Bessou and Damien Cougnaud from the French 'Réseau de transport d'Electricité' (RTE), who greatly helped in making available all the needed GIS information on the French high voltage overhead line network; and (4) all the NRCH research assistants who collected the cases' addresses and the paediatric oncology teams for their help in data collection.

DISCLAIMER

A written contract between RTE and Inserm U1018 team 6 has been concluded and states that team 6 has complete control over the conduct, interpretation and publication of the study. This paper has not been approved by any RTE personnel member other than François Deschamps, who approved it in his capacity as author. The paper does not necessarily represent RTE's views.

REFERENCES

- Ahlbom A, Day N, Feychting M, Roman E, Skinner J, Dockerty J, Linet M, McBride M, Michaelis J, Olsen JH, Tynes T, Verkasalo PK (2000) A pooled analysis of magnetic fields and childhood leukaemia. *Br J Cancer* 83: 692–698.
- Amigou A, Sermage-Faure C, Orsi L, Leverger G, Baruchel A, Bertrand Y, Nelken B, Robert A, Michel G, Margueritte G, Perel Y, Mechinaud F,



- Bordigoni P, Hémon D, Clavel J (2011) Road traffic and childhood leukemia: the escale study (sfce). *Environ Health Perspect* 119: 566–572.
- Bessou J, Deschamps F, Figueroa L, Cougnaud D (2013) Methods used to estimate residential exposure to 50 Hz magnetic fields from overhead power lines in an epidemiological study in France. *J Radiol Prot* 33(2): 349–365.
- Clavel J, Goubin A, Auclerc MF, Auvrignon A, Waterkeyn C, Patte C, Baruchel A, Leverger G, Nelken B, Philippe N, Sommelet D, Vilmer E, Bellec S, Perrillat-Menegaux F, Hémon D (2004) Incidence of childhood leukaemia and non-hodgkin's lymphoma in france: national registry of childhood leukaemia and lymphoma, 1990–1999. *Eur J Cancer Prev* 13: 97–103.
- Draper G, Vincent T, Kroll ME, Swanson J (2005) Childhood cancer in relation to distance from high voltage power lines in england and wales: a case-control study. *BMJ* 330: 1290.
- Greenland S, Sheppard AR, Kaune WT, Poole C, Kelsh MA (2000) A pooled analysis of magnetic fields, wire codes, and childhood leukemia. childhood leukemia-emf study group. *Epidemiology* 11: 624–634.
- IARC (2002) *Monograph on the Evaluation of Carcinogenic Risks to Humans Non-ionizing Radiation, Part 1: Static and Extremely Low Frequency (elf) Electric and Magnetic Fields/ IARC Working Group on the Evaluation of Carcinogenic Risks to Humans Vol. 80*. IARC press: Lyon, France.
- Kheifets L, Ahlbom A, Crespi CM, Draper G, Hagihara J, Lowenthal RM, Mezei G, Oksuzyan S, Schüz J, Swanson J, Tittarelli A, Vinceti M, Wunsch Filho V (2010) Pooled analysis of recent studies on magnetic fields and childhood leukaemia. *Br J Cancer* 103: 1128–1135.
- Kheifets L, Oksuzyan S (2008) Exposure assessment and other challenges in non-ionizing radiation studies of childhood leukaemia. *Radiat Prot Dosimetry* 132(2): 139–147.
- Kroll ME, Swanson J, Vincent TJ, Draper GJ (2010) Childhood cancer and magnetic fields from high-voltage power lines in england and wales: a case-control study. *Br J Cancer* 103: 1122–1127.
- Lacour B, Guyot-Goubin A, Guissou S, Bellec S, Désandes E, Clavel J (2010) Incidence of childhood cancer in france: national children cancer registries, 2000–2004. *Eur J Cancer Prev* 19: 173–181.
- Maslanyj M, Simpson J, Roman E, Schüz J (2009) Power frequency magnetic fields and risk of childhood leukaemia: misclassification of exposure from the use of the 'distance from power line' exposure surrogate. *Bioelectromagnetics* 30: 183–188.
- Maslanyj MP, Mee TJ, Renew DC, Simpson J, Ansell P, Allen SG, Roman E (2007) Investigation of the sources of residential power frequency magnetic field exposure in the uk childhood cancer study. *J Radiol Prot* 27: 41–58.
- Mezei G, Kheifets L (2006) Selection bias and its implications for case-control studies: A case study of magnetic field exposure and childhood leukaemia. *Int J Epidemiol* 35(2): 397–406.
- Schmiel S, Blettner M (2010) The association between extremely low-frequency electromagnetic fields and childhood leukaemia in epidemiology: enough is enough? *Br J Cancer* 103: 931–932.
- Schüz J, Ahlbom A (2008) Exposure to electromagnetic fields and the risk of childhood leukaemia: a review. *Radiat Prot Dosimetry* 132(2): 202–211.
- Schüz J, Grigat JP, Störmer B, Rippin G, Brinkmann K, Michaelis J (2000) Extremely low frequency magnetic fields in residences in germany. distribution of measurements, comparison of two methods for assessing exposure, and predictors for the occurrence of magnetic fields above background level. *Radiat Environ Biophys* 39: 233–240.
- Sermage-Faure C, Laurier D, Goujon-Bellec S, Chartier M, Guyot-Goubin A, Rudant J, Hémon D, Clavel J (2012) Childhood leukemia around french nuclear power plants-the geocap study, 2002–2007. *Int J Cancer* 131: E769–E780.
- Swanson J (2008) Methods used to calculate exposures in two epidemiological studies of power lines in the UK. *J Radiol Prot* 28: 45–59.
- UKCCS (1999) Exposure to power-frequency magnetic fields and the risk of childhood cancer. UK childhood cancer study investigators. *Lancet* 354: 1925–1931.
- WHO (2007) Extremely low frequency fields environmental health criteria. Monograph no 238. available at http://www.who.int/peh-emf/publications/elf_ehc/en/index.html.
- This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License.